

Press Meeting 1 June 2010

二零一零年六月一日新聞發佈會

Dr Wong Ka-hing
Department of Health
衛生署 黃加慶醫生



HIV – routes of transmission

愛滋病病毒的傳播途徑

- Sexual : heterosexual or homosexual
性接觸:異性或同性
- Blood-borne: injecting drug use, contaminated blood/organ/needles
血液接觸:藥物注射、受污染的血液/器官/針具
- Mother-to-child
母嬰傳染

Window period

空窗期

- The time period between contracting HIV and testing positive for HIV antibody, is usually taken to be 3 months.

從感染了愛滋病病毒至愛滋病病毒抗體測試呈陽性反應的期間，一般視為三個月。

HIV Reporting System in Hong Kong

香港的愛滋病病毒感染個案呈報系統

Major fields of information 主要資料:

- Gender 性別
- Ethnicity (Chinese –Y/N) 族裔
- Age (Adult –Y/N) 年齡組別
- Source of reports 呈報來源
- Progression to AIDS 發病(愛滋病)
- Route of transmission 傳播途徑
- Suspected place of infection 估計感染地區

Since 1984

一九八四年開始運作

Voluntary reporting

自願呈報

Anonymous

不記名

Source of reports 呈報來源:


- Physicians 醫生
- Confirmatory laboratories 確認測試化驗室

1984 - first HIV case diagnosed
一九八四年發現首例愛滋病病毒感染

Annual HIV/AIDS Statistics

香港每年愛滋病病毒感染及愛滋病統計

1984 – March 2010, Hong Kong (N=4544/1125)





Gender of reported HIV infection

感染愛滋病毒人士之性別分佈

1984 – March 2010 (N=4544)

Year 2010 (Jan - Mar)





Year 2010 (Jan - Mar)

Ethnicity of reported HIV infection

感染愛滋病病毒人士之族裔分佈



1984 – March 2010 (N=4544)



Year 年份



Age group and gender of reported HIV infection

感染愛滋病病毒人士之年齡及性別分佈
1984 – March 2010 (N=4544)



Routes of transmission of HIV infection


感染愛滋病病毒人士之傳染途徑
1984 – March 2010 (N=4544)



Route of transmission of HIV infection

香港每年感染愛滋病毒人士之傳染途徑分佈

1984 – 2009, Hong Kong (N=4544)



MSM 男男性接觸者

Heterosexual 異性性接觸


Injecting drug use 注射毒品

Blood contact 輸入血液 / 血製品

Perinatal 母嬰傳播


Undetermined 不詳

Heterosexual transmission 異性性接觸感染個案




No. of reports remains at a stable level
報告數字保持平穩

MSM transmission 男男性接觸感染個案



No. of reports is increasing
報告數字呈上升趨勢

Transmission through injecting drug use 注射毒品感染個案



No. of reports is decreasing
報告數字呈下降趨勢





Source of referral of HIV infection


香港每年感染愛滋病毒人士之轉介來源分佈

1984 – March 2010 (N=4544)

Year 2010 (Jan - Mar)




Sources of HIV reports from heterosexual men



異性接觸感染男士多於以下地方確診 infections in **heterosexual men** are mostly referred from

- 公共醫療機構 Public clinics/hospitals (42% in this quarter)
- 私人醫療機構 Private clinics/hospitals (33% in this quarter)

Sources of HIV reports from MSM




男男性接觸者多於以下地方確診 infections in **MSM** are mostly referred from

- 衛生署愛滋病服務組 AIDS Unit of DH (37% in this quarter)
- 愛滋病服務機構 AIDS service organisations (17% in this quarter)
- 公共醫療機構 Public clinics/hospitals (17% in this quarter)
- 衛生署性病門診 DH's STI clinics (17% in this quarter)


Suspected location of HIV infection

估計愛滋病毒病毒感染地區


2006 – March 2010 (N=1719)



Heterosexual Male 異性性接觸男性 (2006 – Mar 2010)



Men who have sex with men 男男性接觸者 (2006 – Mar 2010)



In first quarter of 2010, 62.4% of the newly reported HIV cases were managed at specialist services

二零一零年第一季，62.4%的新呈報感染個案在專科服務接受治療

Natural course of HIV infection

愛滋病病毒感染的自然進程

- Without treatment, about half of the infected people will progress to AIDS in ten years' time.
若沒有接受治療，約一半受感染人士會於十年內發病成為愛滋病患者。
- AIDS is characterised by development of major clinical complications, e.g. opportunistic infections, cancers, related to the underlying immunosuppression.
愛滋病表示因抵抗力下降，出現了嚴重的併發症，例如伺機性感染、惡性腫瘤。
- HIV alone $\neq$ AIDS ; HIV + AIDS-defining illness = AIDS
愛滋病病毒感染 $\neq$ 愛滋病；
愛滋病病毒感染 + 愛滋病界定疾病 = 愛滋病

HIV alone $\neq$ AIDS ;

AIDS = HIV + AIDS-defining illness

愛滋病病毒感染 $\neq$ 愛滋病；


**愛滋病 = 愛滋病病毒感染 + 愛滋病
界定疾病**

Yearly statistics on AIDS 每年愛滋病統計

1985 – March 2010 (N=1125)

Three commonest AIDS –defining illnesses 三種最常見愛滋病界定疾病

	This quarter 本季	Cumulative 累積
PCP 肺囊蟲肺炎	52.0%	39.0%
Tuberculosis 結核病	26.3%	27.3%
Fungal infections 真菌感染	15.8%	16.9%




Trend of AIDS-defining illnesses

愛滋病界定疾病流行情況

1984 – March 2010

Pneumocystis pneumonia and tuberculosis were the most important AIDS-defining illnesses

肺囊蟲肺炎和結核病是最重要的愛滋病界定疾病





AIDS –defining illnesses

愛滋病界定疾病

1985 – March 2010 (N=1125)

Year 2010 (Jan - Mar)



HIV situation in 1st quarter 2010 at a glance

二零一零年第一季度愛滋病流行情況一覽

HIV infections reported	101
--------------------------------	------------

愛滋病病毒感染呈報個案	
-------------	--

Male/Female 男/女	75/26
-----------------	-------

Chinese/Non-Chinese 華裔/非華裔	65/36
----------------------------	-------

Route of transmission 感染途徑	
----------------------------	--

Heterosexual 異性性接觸	20
--------------------	----

Homosexual 同性性接觸	37
------------------	----

Bisexual 雙性性接觸	4
----------------	---

Injecting drug use 注射毒品	4
-------------------------	---

Blood/blood product infusion 輸入血液 / 血製品	0
---	---

Perinatal 母嬰傳播	1
----------------	---

Undetermined 不詳	35
-----------------	----

AIDS reported	19
----------------------	-----------

愛滋病呈報個案	
---------	--

Summary 撮要

- 101 HIV infections were reported to DH in 1st quarter 2010.
二零一零年第一季衛生署共收到一百零一宗愛滋病病毒感染個案。
- Sexual contact was the major route of transmission this quarter. Number of HIV report through homosexual or bisexual transmission increased in recent years. Most of them were infected locally.
性接觸是今季主要的愛滋病病毒傳播途徑。男男性接觸的個案近年增多，大部份個案為本地感染。
- 19 AIDS reports were received in this quarter, a stable level in the past few years.
本季有十九宗愛滋病呈報，數字維持穩定水平。
- *Pneumocystis pneumonia* is the commonest AIDS-defining illness in this quarter.
肺囊蟲肺炎是本季最常見的愛滋病界定疾病。

KEY MESSAGES

重要訊息

1

Anyone with suspected risk should go for an HIV test early

擔心受感染人士應儘快接受愛滋病病毒抗體測試

AIDS Hotline for HIV Testing: 2780 2211

Free, Anonymous and Confidential

愛滋熱線，預約愛滋病病毒抗體測試：

二七八零二二一一

免費、不記名、保密

Social Hygiene Clinics, Department of Health

衛生署社會衛生科診所

Private doctors

私家醫生

Other HIV testing services in Hong Kong

香港其他愛滋病測試服務

-  **AIDS Concern**
關懷愛滋
Tel: 電話: 2898 4411
-  **Hong Kong AIDS Foundation**
香港愛滋病基金會
Tel: 電話: 2513 0513
-  **CHOICE**
再思社區健康組織
Tel: 電話: 3188 9024
-  **SRACP**
香港善導會
Tel: 電話: 2323 3983

KEY MESSAGES

重要訊息

- 2 Use a condom for safer sex**
採取預防措施，正確使用安全套
- 3 Break the needle habit; Methadone does it**
要美沙酮，不要針筒
- 4 Pregnant women should attend early for antenatal HIV testing**
懷孕婦女應儘早作產前愛滋病病毒抗體測試

H
I >
H I V
A I D S
H I V H I
A I D S A I
H I A I D S
I V V I D S H I V A
V A I D S H I A A I
U S M I V V I D S H
< V - D S M I > V I D

要**正視** 要**測試**

Face it! **Test** it!

愛滋熱線 **2780 2211**
AIDS Hotline

及早接受愛滋病毒抗體測試，儘早診斷和治療，繼續健康人生。
Get an HIV Antibody Test without delay. Early checkup, early treatment for healthy life.



相關網址 Related Websites:
<http://www.aids.gov.hk>
<http://www.27802211.com>

<http://www.aids.gov.hk>