

HIV SURVEILLANCE REPORT – 2013 UPDATE

**Special Preventive Programme
Centre for Health Protection
Department of Health
Hong Kong Special Administrative Region
December 2014**

This report is produced and published by:

Surveillance Team
Special Preventive Programme
Centre for Health Protection
Department of Health
Hong Kong Special Administrative Region

Team members:

Dr KH WONG
Dr BY SHU
Dr Philip WONG
Ms WK CHAN
Mr Steve Yim

Correspondence:

Special Preventive Programme
c/o 2/F, Wang Tau Hom Jockey Club Clinic
200 Junction Road East
Kowloon, Hong Kong
Tel: (852) 3143 7225 Fax: (852) 2780 9580
Website: <http://www.aids.gov.hk>
E-mail: aids@dh.gov.hk

CONTENTS

CONTENTS	3
PREFACE	5
ACKNOWLEDGEMENTS	6
ABBREVIATION	7
1. SUMMARY REVIEW	8
Background	8
HIV/AIDS reporting system	9
Discussion	21
2. TABULATED RESULTS OF HIV/AIDS REPORTING	23
System description	23
Box 2.1 Annual and cumulative reports of HIV/AIDS cases	25
Box 2.2 Source of reporting of HIV/AIDS cases	26
Box 2.3 Ethnicity & gender of reported HIV/AIDS cases	28
Box 2.4 Age distribution of reported HIV/AIDS cases	30
Box 2.5 Exposure category of reported HIV/AIDS case	36
Box 2.6 Reported HIV/AIDS cases in injecting drug users (1994– 2013)	38
Box 2.7 Reported sexually acquired HIV/AIDS cases (1994– 2013)	40
Box 2.8 Profile of primary AIDS defining illnesses (ADI) (1994 - 2013)	43
3. TABULATED RESULTS OF HIV PREVALENCE SURVEYS	44
System description	44
Box 3.1 HIV prevalence in blood donors at Hong Kong Red Cross Blood Transfusion Service	46
Box 3.2 HIV prevalence in clients attending Social Hygiene Services, from voluntary blood testing (2004 – 2013)	48
Box 3.3 HIV prevalence in drug users attending methadone clinics	49
Box 3.4 HIV prevalence in drug users attending inpatient drug treatment centres / institutions, from unlinked anonymous screening (2004 - 2013)	49
Box 3.5 HIV prevalence in newly admitted prisoners from unlinked anonymous screening (2004 - 2013)	51
Box 3.6 HIV prevalence in patients attending government TB & Chest Clinics, from voluntary blood testing (2004 - 2013)	52
Box 3.7 HIV prevalence among antenatal women from Universal Antenatal HIV Antibody Testing Programme (2004 - 2013)	53
Box 3.8 HIV prevalence among MSM tested by AIDS Concern (2004 - 2013)	54
Box 3.9 HIV prevalence among MSM – PRISM* (2006, 2009 and 2011)	55

Box 3.10 HIV prevalence among Female Sex Workers – CRISP* (2006 and 2009), HARIS ** (2013)	56
4. TABULATED RESULTS OF STATISTICS ON SEXUALLY TRANSMITTED INFECTIONS (STI)	57
System description	57
Box 4.1 Total number of STI newly reported by individual Social Hygiene Clinic	58
Box 4.2 Annual newly reported STIs in Social Hygiene Clinics	60
Box 4.3 Syphilis newly reported by Social Hygiene Clinics (2009 - 2013)	61
Box 4.4 Sexually acquired HIV infection in Hong Kong (1994-2013)	62
Box 4.5 Syndromic presentations of STI from Behavioural Survey of Social Hygiene Service (2009-2013)	63
5. TABULATED RESULTS ON BEHAVIOURAL MONITORING	64
System description	64
Box 5.1 Median number of sex partners in the previous year among adult ^ heterosexual men / MSM attending AIDS Counselling and Testing Service (ACTS) (2004-2013)	66
Box 5.2 Recent history* of commercial / casual sex among adult ^ heterosexual men (2004-2013)	67
Box 5.3 Condom use with regular partners among adult heterosexual men	68
Box 5.4 Condom use with commercial / casual partners among adult heterosexual men	70
Box 5.5 Condom use among Men have Sex with Men (MSM)	72
Box 5.6 Proportion of current injectors* (2004-2013)	74
Box 5.7 Proportion of current needle-sharers* (2004-2013)	75
6. TABULATED RESULTS OF HIV-1 GENOTYPING STUDIES	76
System description	76
Box 6.1 Proportion of reports* with subtypes by year in Hong Kong, 2001 - 2013	77
Box 6.2 Distribution of HIV-1* subtypes	78
Box 6.3 Trend in the common HIV-1* subtypes in Hong Kong, 2001 – 2013	79
Box 6.4 Trend in HIV-1* subtype CRF01_AE in Hong Kong, 2001 – 2013	80
Box 6.5 Trend in HIV-1* subtype B in Hong Kong, 2001 – 2013	83
Box 6.6 Trend in HIV-1* subtype C in Hong Kong, 2001 – 2013	86
Box 6.7 Prevalence of intermediate or high level drug resistance related mutation among newly diagnosed HIV patients, 2003-2012	89
APPENDIX I: HIV/AIDS REPORT FORM (DH2293)	90
APPENDIX II: CLASSIFICATION SYSTEM FOR HIV INFECTION AND SURVEILLANCE CASE DEFINITION FOR AIDS IN ADOLESCENTS AND ADULTS IN HONG KONG.	91
APPENDIX III: CONDOM DISTRIBUTION FOR THE PREVENTION OF HIV AND STI BY DEPARTMENT OF HEALTH	92

PREFACE

The increasing trend of annual number of newly reported HIV infection continued in 2013 and reached a record high of 559 cases. Sexual transmission remained the major route of HIV transmission in Hong Kong in 2013, while the transmission from other routes including drug injection have been kept at a relatively low level. Overall, Hong Kong is still enjoying a low prevalence level of HIV infection in the general population.

Similar to other developed countries, Hong Kong is facing a continuous threat of increasing HIV infection in the men who have sex with men (MSM) community in recent years. The HIV prevalence among local MSM was still the highest amongst all at risk populations. In addition, a potential upsurge of infection among injecting drug users (IDU) is still a concern from the worldwide and regional experience on HIV and drug.

With the expansion of community-based HIV voluntary testing services, non-governmental organisations were playing an increasing role in understanding the local HIV epidemiology especially among the at-risk populations including MSM, IDU and Female Sex Workers (FSW). Many non-governmental organisations are conducting HIV prevalence & behavioral surveys in different at-risk populations through their service networks.

This *annual surveillance report* is an initiative of Special Preventive Programme (SPP), Centre for Health Protection, Department of Health. The report aims to provide strategic information to facilitate planning of services and intervention activities for the prevention, care and control of HIV/AIDS. Following a commentary, data collected from five main components of our surveillance programme (the HIV/AIDS voluntary reporting system, HIV prevalence surveys, Sexually Transmitted Infections caseload statistics, behaviour studies and HIV-1 genotyping studies) are presented as tables and graphs. Some changes have been made in this year's report to enhance its contents including that selected findings of the HIV and AIDS Response Indicator Survey (HARIS) were described.

Electronic copy of this report is accessible in our website <http://www.aids.gov.hk>. Moreover, the quarterly bulletins, factsheets on yearly situation and specific surveys, and other information relating to HIV surveillance and epidemiology are also available in the website. Your comments and suggestions are always welcome.

Surveillance team
Special Preventive Programme
Centre for Health Protection
Department of Health
December 2014

ACKNOWLEDGEMENTS

This report is only achievable with the concerted efforts contributed by many different stakeholders. First and foremost, we would like to thank our colleagues of the Social Hygiene Service, the Narcotics and Drug Administration Unit, Tuberculosis & Chest Service, Family Health Service, Surveillance and Epidemiology Branch and the Virology Division of Public Health Laboratory Services Branch who have provided the necessary data and support over the years. For data collected in the prison setting, we are indebted to the staff of the Correctional Services Department for their invaluable assistance in carrying out HIV prevalence surveys on a regular basis.

Secondly, special thanks are dedicated to many agencies that have helped collect and update the relevant statistics referred by this report. They included the Hong Kong Red Cross Blood Transfusion Service, the Society for the Aid and Rehabilitation of Drug Abusers, AIDS Concern, the Narcotics Division of the Security Bureau, the Department of Microbiology of the University of Hong Kong, the School of Public Health and Primary Care of the Chinese University of Hong Kong, many of our local AIDS and non-AIDS non-governmental organisations and various public hospitals/clinics, in particular Queen Elizabeth Hospital, Prince of Wales Hospital and Princess Margaret Hospital. We also take this opportunity to thank all physicians, health care professionals and related workers who have contributed to HIV/AIDS reporting and other surveillance components.

Finally, we must thank the usual excellent support from the SPP staff in terms of collecting, collating, compiling and analyzing the information as well as the editing and production of this report.

ABBREVIATION

ACTS	AIDS Counseling and Testing Service
ADI	AIDS Defining Illness
AIDS	Acquired Immune Deficiency Syndrome
AC	AIDS Concern
AIMSS	Asia Internet MSM Sex Survey
CDC	Centers for Disease Control and Prevention
CRISP	Community based Risk behavioral and SeroPrevalence survey for female sex workers
CD4	Cluster of Differentiation (CD) 4 molecule
CHOICE	Community Health Organisation for Intervention, Care and Empowerment
CRDA	Central Registry of Drug Abuse
CHP	Centre for Health Protection
CRF	Circulating Recombinant Form
DH	Department of Health
DRS-M	Drug Rehabilitation Services – Methadone clinics
DRS-S	Drug Rehabilitation Services – Shek Kwu Chau Treatment and Rehabilitation Centre
ELISA	Enzyme-linked Immunosorbent Assay
FSW	Female Sex Worker
HE	Heterosexual
HAART	Highly Active Antiretroviral Therapy
HARIS	HIV and AIDS Response Indicator Survey
HIV	Human Immunodeficiency Virus
IDU	Injecting Drug User
ITC	Integrated Treatment Centre
MUT	Methadone Universal HIV Antibody (Urine) Testing
MSM	Men who have Sex with Men
NSGI	Non-specific Genital Infection
NGU	Non-gonococcal Urethritis
PCP	Pneumocystis Pneumonia
PCR	Polymerase Chain Reaction
PRISM	HIV Prevalence and Risk behavioral Survey of Men who have sex with men
SADRA	The Society for the Aid and Rehabilitation of Drug Abusers
SKC	Shek Kwu Chau Treatment and Rehabilitation Centre
STI	Sexually Transmitted Infection
SPP	Special Preventive Programme
SHS	Social Hygiene Service
SAS	Street Addict Survey
TB	Tuberculosis
ul	microliter

1. SUMMARY REVIEW

Background

1. The HIV surveillance system in Hong Kong comprises 5 main programmes to provide a detailed description of the local HIV/AIDS situation. They are (a) voluntary HIV/AIDS case-based reporting; (b) HIV prevalence surveys; (c) sexually transmitted infections (STI) caseload statistics; (d) behavioral studies; and (e) HIV-1 genotyping studies. The data is collected, analyzed and disseminated regularly by the surveillance team of Special Preventive Programme (SPP), Centre for Health Protection (CHP), Department of Health (DH). At present, the latest HIV/AIDS statistics are released at quarterly intervals at press media briefings and in electronic format (<http://www.aids.gov.hk>). Data from various sources are compiled annually and released in this report.

2. The following paragraphs highlight the main findings from HIV/AIDS surveillance activities undertaken in 2013 and before. Please refer to the following pages for the details of the programmes.

HIV/AIDS reporting system

3. The Department of Health has implemented a voluntary anonymous case-based HIV/AIDS reporting system since 1984, which receives reports from doctors and laboratories. Doctors report newly diagnosed HIV cases by a standard form (DH2293) which was lately revised in March 2010 with the data field on “date of last negative HIV test” added. Before 2006, only cases with Western Blot confirmed HIV antibody positive laboratory result were counted as HIV infection for cases aged above 18 months. With increase detection, those cases with PCR positive result and clinical or laboratory indication of recent infection were also counted as confirmed HIV infection in the reporting system since the 4th quarter of 2006.

4. In 2013, DH received 559 HIV and 84 AIDS reports (Box 2.1). The number of HIV cases in 2013 reached a record yearly high and continued the increasing trend since 2011, after the slight decrease in 2009 and 2010. This brought the cumulative total to 6342 and 1437 for HIV and AIDS reports respectively. Under the revised definition, 11 PCR positive cases with clinical or laboratory indication of recent infections were included as HIV infection in 2013. Public hospitals/clinics/laboratories were still the commonest source of HIV reports in 2013, which accounted for 34.9% of all. Private hospitals/clinics/laboratories and Social Hygiene Clinics were other common sources of HIV reports, which account for 17.7% and 15.7% respectively. (Box 2.2)

5. In 2013, around 79% of reported HIV cases were male. The male-to-female ratio was 3.9:1 in 2013, which remained relatively stable as compare to that of 3.5:1 in 2012. About 70% of reported cases were Chinese. Asian non-Chinese accounted for 9% of reports. (Box 2.3) The median age of reported HIV cases was 36 (Box 2.4) and 20-49 was the commonest age group in both male and female cases. There was one children (age <13) HIV case reported in 2013. Around 80% of reported HIV cases were believed to have acquired the virus through sexual transmission in 2013, including homosexual (50%), heterosexual (26%), and bisexual exposure (4%). Injecting drug use accounted for 1% of reported HIV infections. There was 1 case of HIV transmission via blood/blood product which occurred outside Hong Kong and no case of infection via perinatal route in 2013. The suspected routes of transmission were undetermined in around one-fifth (19%) of cases. This means that, after excluding those with undetermined exposure category, sexual transmission accounted for about 98% among HIV reports with defined risks. (Box 2.5(a))

HIV Surveillance at a glance (2013)


- 559 HIV reports and 84 AIDS reports
- Gender: 79.4% male
- Ethnicity: 70.1% Chinese
- Age: Median 36
- Risks:
 - 53.7% Homosexual/bisexual contact
 - 26.3% Heterosexual contact
 - 1.2% Injecting drug use
 - 0.2% Blood transfusion
 - 18.6% Undetermined
- CD4 at reporting: Median 286/ul
- HIV-1 subtypes: commonest are CRF01_AE and B
- Commonest primary AIDS defining illness: PCP and TB
- HIV prevalence
 - Blood donors: <0.01%
 - Antenatal women: 0.01%
 - STI clinic attendees: 0.34%
 - Methadone clinic attendees: 0.68%

The rising trend in men who have sex with men (MSM) cases continued

6. Similar as previous few years, sexual contact including both heterosexual and homosexual/bisexual, remained the commonest route of HIV transmission in Hong Kong in 2013, which accounted for 80%. In the early years of HIV/AIDS epidemic in Hong Kong around 1980s and early 1990s, it used to report more cases from men who have sex with men, who had homosexual or bisexual contacts. The trend then reversed with heterosexual transmission overtaking homosexual / bisexual transmission from 1993 onwards. Since 2004, a rising trend in MSM has been observed again and the situation continued in 2013 with 300 MSM cases (65.9%) identified out of 455 cases with defined risks. (Box 2.5(a)).


7. The high weighting of MSM among male HIV cases remained in 2013. 67.6% of male HIV reports in 2013 contracted the virus through homosexual or bisexual contact. Heterosexual contact in male cases accounted for about 16.2%, whereas the routes of transmission were undetermined in another 14.6% of the male cases. The ratio of heterosexual men against MSM gradually dropped from its peak of 4.2:1 in 1998 to 0.8: 1 in 2005 and further to 0.2:1 in 2013. (Box 1.1 and 2.7(c)) These showed the increasing proportion of MSM among the male infected cases. Similar trend of increasing AIDS cases among MSM was observed, the ratio of heterosexual men against MSM decreased from 23.5:1 in 2000 to 0.4:1 in 2013.

Box 1.1 The number of MSM cases has taken over heterosexual men cases in the reporting system since 2005 and the gap continued to widen.




8. In 2013, majority of the MSM cases were Chinese (84.3%) and of age group 20-49 (91.3%). A rising trend in the number of reported Chinese MSM cases was observed in recent years despite a modest drop between 2007 and 2008. (Box 1.2) In 2013, the median age of MSM cases at report was 33, which was lower than 43.5 of heterosexual male cases. The median age of HIV infected MSM population, has shown a decreasing trend in the past few years from 37 in 2010 to 34 in 2012 and 33 in 2013. (Box 1.3) In 2013, age group 20-29 was the commonest age group of reporting in MSM, which accounted for 34.0%, followed by age group 30-39 (33.3%) and age group 40-49 (24.0%). (Box 1.4) Reported data since 2006 suggested that a relatively higher proportion of MSM infections were occurred in Hong Kong, as compared to a much lower proportion in heterosexual men. In 2013, around 75.3% of MSM infection occurred in Hong Kong while only around 50% of local heterosexual male infection. (Box 1.5)


Box 1.2 Ethnicity Breakdown of HIV-infected MSM cases (2004-2013)



Box 1.3 Median HIV reporting age of HIV-infected MSM cases, heterosexual man and heterosexual women (2004-2013)




Box 1.4 Age breakdown of HIV-infected MSM cases (2004 - 2013)




Box 1.5 Suspected location of HIV cases (2006 - 2013)

(a) MSM



(b) Heterosexual men



HIV prevalence among men who have sex with men persistently higher than other at-risk population

9. The third community-based survey (PRiSM) in gay saunas, bars and beaches conducted in 2011 revealed that the HIV prevalence among local MSM was around 4.08%, which remained relatively stable as compared to 4.05% and 4.31% in the previous two rounds in 2006 and 2008 respectively. (Box 1.6 and Box 3.9) The prevalence among MSM was persistently higher than other at-risk population such as female sex worker (Box 3.10) and drug users (Box 3.3 and Box 3.4).

10. A subproject iPRiSM which involved the conduction of the survey through internet was also added in the 2011 round for the first time, in addition to the traditional venue-based surveillance. The HIV prevalence was around 3.3%, as compared to a self-reported 5.5% in an on-line survey Asia Internet MSM sex survey (AIMSS) conducted in 2010.

11. AIDS Concern's voluntary HIV testing service targeting MSM was another data source to estimate the HIV prevalence in the local MSM community, despite that the data may be affected by participant bias to a certain extent. It showed a prevalence of 1.81% in 2013 which remained relatively stable in the past few years. (Box 3.8) The HIV prevalence among MSM remained at a relatively stable level of around 2% in the past few years.

Condom use and HIV testing among men who have sex with men showed an increasing trend

12. The first HIV and AIDS Response Indicator Survey (HARiS) for MSM was conducted in 2013, it showed that the condom use rate in the last anal sex with emotional relationship partner, regular sex partner and non-regular sex partner were 63.7%, 76.7% and 79.5% respectively. (Box 1.6) The condom use rate during last anal sex with regular sex partner showed an increasing trend as compared to the latest round of PRISM conducted in 2011, which was 61.9%. The condom use rate during last anal sex with non-regular sex partner remained relatively stable at around 80% in the past few years.

13. Both the ever HIV testing rate (74%) and HIV testing rate in past one year (57%) increased in the 2013 HARiS, as compared with the 2011 PRISM findings (67% and 40% respectively). It might suggest an increased awareness to undergo HIV testing and even regular testing in the MSM community.

Box 1.6 Comparison between 2011 PRISM and 2013 HARiS results

Results	PRISM 2011		HARiS 2013 venue-based, centre-based and internet based
	Venue-based	Internet-based	
Sample Size	816	180	853
Adjusted HIV prevalence	4.08% (95% CI 3.44-4.85%)	3.3% (95% CI 1.54-7.08)	/
Condom use in last anal sex with:			
-emotional relationship partner (ESP)	/	/	63.7%
-Regular Sex Partner (RSP)	61.9%	60.0%	76.7%

-Non-regular Sex Partner (NRSP)	82.7% (inside HK) 81.2% (outside HK)	81.4% (inside HK) 79.2% (outside HK)	79.5%
-Commercial Sex Partner (CSP)	/	/	69.9%
HIV testing			
-Ever test for HIV	67%	63%	73.7%
-HIV test within past year	40%	41%	57.0%

14. According to the survey conducted among the clients attending the DH's AIDS Counseling and Testing Service (ACTS), the median number of casual sex partners among MSM was consistently higher than those heterosexual men, 3 in 2013. (Box 5.1) The consistent condom use rate among MSM with regular partners and casual partners remained stable in 2013, which was 47.3% and 56.9% respectively, as compared with the rate of 45.3% and 59.3% respectively in 2012. (Box 5.5(a)) On the contrast, the condom use rate for last anal sex with both regular partners and casual partners (59% and 64.9% respectively) showed a slight decreasing trend in 2013, as compared with 60.6% and 75.8% respectively (Box 5.5(b)).

15. Additional behavioural data derived from MSM attending AIDS Concern's testing service showed that the consistent condom use rate for boyfriend, regular sex partners and casual sex partners in 2013 was 47.7%, 59.6% and 67.2% respectively. (Box 5.5)

The proportion of heterosexual cases remained stable

16. In 2013, there was a total of 147 heterosexual cases reported, which accounted for about one-quarter of the all reported HIV cases, and was similar to the proportion in 2012. (Box 2.5(a)) The proportion of heterosexual cases among all reported HIV cases gradually dropped from its peak of 71% in 1998 to 37% in 2005 and 26% in 2013. The female to male ratio for heterosexual cases gradually increased in the past decade from 0.5:1 in 2004 to 1:1 in 2013, which showed increasing female proportion in heterosexual cases. The median age of heterosexual cases in 2013 was 36 for female and 43.5 for male respectively. In 2013, heterosexual male cases were mainly Chinese (76%) whereas Chinese accounted for around half (52%) of female heterosexual cases.

17. STI caseload statistic from Social Hygiene Clinics is an important component of the local HIV surveillance programme as the presence of STI is an indicator of high risk sexual behaviors which also increase the risk of contracting HIV. In 2013, 15.7% of reported cases was referred from Social Hygiene Clinics. The consistent condom use rate among heterosexual men attending Social Hygiene Clinics with commercial / casual partners in the past 3 months in 2013 was 49.1%, which slightly decreased as compared with 52.9% in

2012. (Box 5.4) This condom use rate remained at only around 50% in the past years, which is an area that needs to be addressed (Box 5.4(a)). Moreover, more than one third of the STI cases were without any symptoms which may delay the diagnosis and the link to appropriate medical care. (Box 4.5). The HIV prevalence of Social Hygiene Clinic attendees remained stable in the previous few years at around 0.2% (0.34% in 2013). (Box 3.2) The total number of STI cases in Social Hygiene Clinics remained stable in the past few years, with an aggregate of 12,912 cases in 2013. (Box 4.1, 4.2)

18. The consistent condom use level observed among those attending AIDS Counseling and Testing Service (ACTS) slightly decreased from 76.6% in 2012 to 74.9 % in 2013 for commercial partners and maintained at 60.9% in 2013 for commercial/causal partners. Discrepancy was noticed when the condom use rate from client's side was compared with that from the sex worker's side. In the HIV and AIDS Response Indicator Survey (HARIS) for female sex worker conducted in 2013, a relatively higher condom use level was revealed among female sex workers in Hong Kong, that the condom use rate in the last intercourse with their regular clients and casual clients was 97.2% and 99.0% respectively.

New HIV infection among drug users remained low but significant level of risky behaviors reported

19. In 2013, the reporting system recorded 7 cases of HIV transmission through injecting drug use, which accounted for 1.2% of all reported cases. The number continued to show a decreasing trend from the peak of 58 cases in 2006 to 14 cases in 2011 and 7 cases in 2013. (Box 2.5(a)) All of the cases in 2013 were male and majority were Chinese (86%). (Box 2.6(a)) The median age was 40. 2 out of the 7 injecting drug user cases were reported from methadone clinics.

20. The Methadone Universal HIV Antibody (Urine) Testing Programme (MUT) launched in 2004 replaced the unlinked anonymous screening (UAS) in methadone clinics to enhance HIV surveillance as well as individual diagnosis and subsequent care of the infected methadone clients. Among those 9674 methadone clinic attendees in 2013, 6908 clients have been tested for HIV with an overall HIV coverage rate of 71.4%. Forty seven clients were found to be positive for HIV and the overall HIV prevalence of methadone clinic attendees in 2013 was 0.68%. (Box 3.3)

21. The proportion of drug users who were currently injecting drugs ranged from about 24% to 86% across different surveys in 2013. (Box 5.6) Moreover, various surveys revealed that around 2% to 24% of the current drug injectors were still practicing needle sharing behaviours, which posed them to the risk of contracting HIV. (Box 5.7) As such, despite that reported HIV infection cases among injecting drug users remained at a low level in 2013, the potential risk of outbreak of HIV among drug users cannot be neglected.

One case of transmission via blood/blood product transfusion recorded

22. In 2013, there was 1 reported case of HIV infection via contaminated blood or blood product transfusion, which occurred outside Hong Kong. The HIV prevalence of new blood donors at Hong Kong Red Cross Blood Transfusion Service remained at a low level of 0.002% in 2013 (Box 3.1(b)).

23. In 2013, there was no perinatal transmission case reported. Since the launch of the Universal Antenatal HIV Testing in September 2001, around 40,000-50,000 pregnant women attending public antenatal services were tested for HIV every year. The coverage of the programme remained at a high level (98.5% in 2013) and the prevalence of HIV infection in pregnant women was found to be stable at around 0.01% in the previous years (0.01% in 2013). Seven pregnant women were tested positive in the programme in 2013. (Box 3.7) Four women subsequently delivered their babies by Caesarean Sections, one by vaginal delivery, one case without delivery mode information, while remaining one cases was without sufficient information. Of these six newborn babies, five were put on anti-retroviral chemoprophylaxis. None of the babies was confirmed to have HIV infection at the time of report writing.

Reconstruction of risk factor for cases without reported route of transmission


24. As the HIV/AIDS case-based reporting system in Hong Kong is entirely voluntary and anonymous, the completeness of the local surveillance database depends largely on the percentage of cases with the report form DH2293 received from attending physicians. Incomplete data due to cases without a risk factor reported may pose a risk of skewing the local epidemic picture. In 2013, 19% of the infected cases did not have a suspected route of transmission reported, as compared to around 21% in 2012. (Box 2.5(a)) A systematic reconstruction method proposed by Dr. Tim Brown was used since 2010 in order to factor in the weightings of undetermined risk cases, to assess the risk for local transmission and to plan and guide appropriate preventive actions.

25. Reconstruction was carried out by assigning one suitable transmission to the undetermined cases. After the analysis of the features of these cases with undetermined risk factor and the prevailing epidemic, it was assessed that all female infections shall be assumed to be heterosexual transmission, unless there is clear indication suggesting otherwise. As for the male cases of undetermined risk factor, it was assessed that they shall be assumed to be either heterosexual contact or homosexual contacts as the risk factor of transmission, subject to the observed ratio in the prevailing year between heterosexual and homosexual contact, providing there is no other indication suggesting otherwise.


26. By using the above methodology of reconstruction, a modified epidemic was constructed by applying our local 10-year data from 2004 to 2013. (Box 1.7(a) and Box 1.7(b)). After the reconstruction, the cases of MSM and heterosexual female showed a mark increase since 2007, while the change in heterosexual male appeared to be relatively moderate. (Box. 1.7 (c)). Although this method might simplify the complex local epidemic, it provides one possible solution to fill the gap in the HIV surveillance system information.

Measures to promote the return rate of report forms from physicians regarding each HIV case have also been implemented in the past few years.


Box 1.7(a) HIV reports before risk factor reconstruction (2004-2013)



Box 1.7(b) HIV reports after risk factor reconstruction (2004-2013)



Box 1.7(c) HIV reports before and after risk factor reconstruction in MSM, heterosexual male and heterosexual female cases (2004-2013)



Regular HIV testing before diagnosis was still not a norm in Hong Kong

27. The HIV/AIDS Report Form (DH2293) was lately revised in March 2010 and become available for reporting use since July 2010 with one data field added to capture the previously negative HIV result among the newly diagnosed cases. The data helps to inform the epidemiology of those cases who were recently infected. Among the 559 cases reported in 2013, data of the HIV/AIDS Report Form was available in 482 cases, of which only 201 cases (41.7%) had the data on previously negative HIV results, which implied regular testing among HIV patients before their diagnoses was rare. Among those 201 cases, seventy-four (36.8%) had previously negative HIV results within one year of the HIV diagnosis, i.e. recently HIV seroconvert and this suggested that at least one-third of the cases were recently infected. However, it was not possible to judge whether the cases with previously negative HIV results beyond one year of HIV diagnosis were recently HIV seroconvert or not, as the observation was limited by the infrequent testing behaviour.

Pneumocystis Pneumonia and Tuberculosis remained the commonest Primary AIDS Defining Illnesses

28. Since the introduction of highly active antiretroviral therapy (HAART) in Hong Kong around 1997, the annual number of reported AIDS cases has been dropping since then and remained at a relatively stable level of around 80 cases per year in the past decade. A total of 84 AIDS cases were reported in 2013 as compared with 86 cases in 2012 (Box 2.5(b)). Majority (89%) of the AIDS reports in 2013 had their AIDS diagnosis within 3 months of HIV diagnosis, suggesting late presentation of the cases.

29. *Pneumocystis jirovecii* pneumonia (previously named *Pneumocystis carinii*) was the commonest ADI in Hong Kong in 2013 which accounted for 44% (37 cases), which is similar to the proportion in 2012. The second most common primary ADI reported in 2013 was *Mycobacterium tuberculosis* which accounted for 20% (17 cases). They were followed by other fungal infections (12%), *Cytomegalovirus* diseases (5%) and *penicilliosis* (4%). (Box 2.8) The university voluntary testing has literally replaced unlinked anonymous screening at TB & Chest Clinics since 2009 in informing the HIV prevalence among TB patients. In 2013, the HIV testing coverage in patients attending government TB & Chest Clinic was 88.2% and HIV prevalence was 0.68%, which remained at a low level of less than 1% in the past few years. (Box 3.6)

Median CD4 of newly reported HIV cases showed an increasing trend but those of older patients remained at a relatively lower level

30. The median CD4 of newly reported HIV cases in 2013 was 286/ul, which was higher than previous few years suggesting that more cases were diagnosed at a relatively earlier stage. The proportion with CD4 \geq 200/ul in 2013 was 63.8%, which was similar to the percentage in previous years. Reporting of CD4 level was also becoming a routine practice in physician, which provided useful information on the timing of diagnosis in the course of HIV infection. In 2013, 78.2% of HIV cases had their CD4 level at diagnosis reported, which was higher than that in the past few years. (Box 1.8) The median CD4 for those aged less than 55 was 309/ul in 2013, which has increased as compared to 300/ul in 2012. On the contrast, the median CD4 count among those who are aged 55 or above has decreased from 197.5/ul in 2012 to 111.5% in 2013. It was consistently lower than the younger group, suggesting that more patients reported at age 55 or above were diagnosed at a relatively late disease stage. (Box 1.9)

Box 1.8 – Reported CD4 levels at HIV diagnosis

Year	No. of HIV reports	No. of CD4 reports (%)	Median CD4 (cell/ul)	CD4 \geq 200 (cell/ul) (%)
2004	268	183 (68.3%)	208	97 (53.0%)
2005	313	239 (76.4%)	201	120 (50.2%)
2006	373	298 (79.9%)	233.5	163 (54.7%)
2007	414	327 (79.0%)	236	182 (55.7%)
2008	435	315 (72.4%)	193	154 (48.9%)
2009	396	289 (73.0%)	278	181 (62.6%)
2010	389	291 (74.8%)	208	149 (51.2%)
2011	438	312 (71.2%)	258	185 (59.3%)
2012	513	386 (75.2%)	280.5	251 (65.0%)
2013	559	437 (78.2%)	286	279 (63.8%)

Box 1.9 – CD4 Reports by age group*

Age	Year	No. of HIV reports	No. of CD4 reports (%)	Median CD4 (cell/ul)	% of CD4 ≥ 200 (cell/ul)
<55	2004	225	161 (71.6%)	218	(55.3%)
	2005	282	216 (76.6%)	199.5	(50.0%)
	2006	341	272 (79.8%)	243.5	(57.4%)
	2007	377	300 (79.6%)	249	(57.3%)
	2008	380	272 (71.6%)	217	(52.6%)
	2009	357	260 (72.8%)	296.5	(66.5%)
	2010	353	259 (73.4%)	219	(52.5%)
	2011	384	275 (71.6%)	280	(62.2%)
	2012	463	346 (74.7%)	300	(66.8%)
	2013	501	387 (77.2%)	309	(68.0%)
≥55	2004	32	22 (68.8%)	93	(36.4%)
	2005	29	23 (79.3%)	223	(52.2%)
	2006	29	26 (89.7%)	154.5	(26.9%)
	2007	33	27 (81.8%)	90	(37.0%)
	2008	53	43 (81.1%)	74	(25.6%)
	2009	38	29 (76.3%)	72	(27.6%)
	2010	36	32 (88.9%)	121	(40.6%)
	2011	53	37 (69.8%)	126	(37.8%)
	2012	48	40 (83.3%)	197.5	(50.0%)
	2013	58	50 (86.2%)	111.5	(32.0%)

*: there may be a slight discrepancy between the sum of individual reports in Box 1.9 and the figures showed in Box 1.8 because of unknown age.

The commonest HIV-1 subtypes were CRF01_AE and B, but genetic diversity continued to increase. The level of drug resistance mutation remained low.

31. In 2013, about 87% of HIV reports had their subtypes documented, at a comparable level as in the past years. (Box 6.1) Subtype CRF01_AE and B of HIV-1 strains remained the first and second most common subtypes identified in Hong Kong, respectively at 44% and 35% of all cases having subtype identified from 2001 to 2013. In 2013, they together accounted for 69% of all HIV cases with subtype documented. (Box 6.2) Over the past decade, CRF_01AE was found to be commoner in female, Asian non-Chinese, heterosexuals and IDU, but became more common in Chinese and male since 2012 (Box 6.4) On the other hand, subtype B was consistently commoner in male, MSM, Chinese and Caucasian. (Box 6.5) Subtype C was commoner in female, Asian non-Chinese and heterosexual (Box 6.6). Over the past few years, both the proportion of Subtype CRF01_AE and B showed a decreasing trend. In contrast, an increasing trend of diversity in subtypes and its circulating recombinant forms was noted, in particular since 2009. (Box 6.3) The proportion of subtype CRF07_BC have increased from 3.4% in 2008 to 8.0% in 2013 while that subtype CRF08_BC increased from 0.8% to 5.8% respectively.

32. According to the HIV resistance threshold survey conducted since 2003, the prevalence of intermediate or high level Drug Resistance related mutations in 2012 was 2.3%, which maintained at a relatively low level in the past few years (from 0% to 4.3%) (Box. 6.7).

Discussion

33. The rising trend of HIV reports continued since 2011 and reached a record high level in 2013, after a modest drop in 2009 and 2010. The total number of HIV reports in 2013 was 559, which increased for about 9% as compared to 513 cases in 2012. The increasing number of MSM reported cases remained the major factor contributing to the high HIV level. The level of heterosexual contact remained stable in the past few years after a peak in 2008 and the number of cases among injecting drug users also remained at a relatively low level of around 10 cases per year since 2009.

34. The number of HIV reports among MSM continued to increase and accounted for the largest proportion of cases in 2013. From the data of previous few years, the increasing trend will likely to be continued in the near future and play a significant role in the local epidemic. Using the reconstruction methodology described in paragraph 25 above, we can observe an ever more dramatic increase in the infection cases among MSM. The latest community-based HIV prevalence survey (PRISM) among MSM in 2011 revealed a HIV prevalence of 4.08%, which was similar to the previous studies in 2006 and 2008 but was considerably higher than those of other risk population including the female sex workers and drug users. As gauged from the PRISM surveys and HARiS survey for MSM 2013, the condom use rate with different types of partners improved over the past years. The HIV testing rate has also increased which may reflect a growing norm of regular HIV testing among MSM community, and could partly explain the continuous increase in the number of new infection in the community. Although majority of the MSM cases (75.3%) were infected locally in 2013, potential threat of HIV infections contracted from neighboring

cities/countries should not be taken lightly due to the increasing cross-border sexual activities in the population.

35. Heterosexual transmission remained relatively stable over the past few years of around 130 cases per year. The proportion of female among heterosexual cases continued to increase and reached a record high of over half (51.0%) in 2013. Upon reconstruction of undetermined female cases, it showed an even more obvious increase for female heterosexual cases. The HIV prevalence in social hygiene clinics attendees and antenatal women remained at a relatively low level in the past decade and was 0.34% and 0.01% in 2013 respectively. However, consistent condom use rates of commercial / casual sex especially gauged from the surveys of heterosexual male remained far from satisfactory and could pose a threat of rebound in the number of cases via heterosexual route.

36. The number of injecting drug cases remained stable of less than 10 cases in the past 2 years (7 cases in 2013). Despite that, the proportion of injection and risky needle-sharing behaviour among the drug users as gauged from several surveys remained at a significant level which poses a potential risk of cluster outbreak and rapid upsurge of infection among the population. This create challenge for both surveillance effort and preventive intervention in the near future. Moreover, the HIV testing coverage in methadone clinics showed a decreasing trend in the past few years which may defer diagnosis and subsequent care of infected drug users.


37. In conclusion, the number of newly reported HIV infections in Hong Kong continued to increase in 2013. Similar to the situation in many developed countries and neighboring areas, MSM infection continued to dominate the HIV epidemic in Hong Kong and will likely to exert its effect in the near future. The situation of heterosexual population and injecting drug user population was relatively stable thus far. Due to the considerable proportion of local reported cases were acquired outside Hong Kong and increasing popularity of cross-border sexual activity, the situation of neighboring cities/countries play an important factor influencing the local HIV epidemiology. In 2013, the HIV prevalence among the general population in Hong Kong was estimated to remain at a low level of about 0.1% and the number of people living with HIV was estimated to be around 4300 based on the projection using Asian Epidemic Model. To combat the HIV epidemic, continuous and collaborative effort in HIV prevention is essential.

2. TABULATED RESULTS OF HIV/AIDS REPORTING


System description

- The HIV/AIDS reporting system is a case-based notification system conducted on a voluntary, anonymous and confidential basis since 1984, with input from physicians and laboratories.

System layout




Box 2.1 Annual and cumulative reports of HIV/AIDS cases




Box 2.2 Source of reporting of HIV/AIDS cases

(a) Year 2013

(i) HIV




(ii) AIDS




(b) Cumulative (1984 - 2013)

(i) HIV



(ii) AIDS



Box 2.3 Ethnicity & gender of reported HIV/AIDS cases

(a) Year 2013

Ethnicity	HIV			AIDS		
	Male	Female	Total	Male	Female	Total
Chinese	345 (77.7%)	47 (40.9%)	392 (70.1%)	59 (84.3%)	7 (50.0%)	66 (78.6%)
Non-Chinese	77 (17.3%)	67 (58.3%)	144 (25.8%)	11 (15.7%)	7 (50.0%)	18 (21.4%)
Asian	27 (6.1%)	25 (21.7%)	52 (9.3%)	7 (10.0%)	4 (28.6%)	11 (13.1%)
White	35 (7.9%)	1 (0.9%)	36 (6.4%)	3 (4.3%)	0 (0.0%)	3 (3.6%)
Black	5 (1.1%)	14 (12.2%)	19 (3.4%)	1 (1.4%)	3 (21.4%)	4 (4.8%)
Others	10 (2.3%)	27 (23.5%)	37 (6.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Unknown	22 (5.0%)	1 (0.9%)	23 (4.1%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Total	444 (100.0%)	115 (100.0%)	559 (100.0%)	70 (100.0%)	14 (100.0%)	84 (100.0%)

(b) Cumulative (1984 - 2013)


Ethnicity	HIV			AIDS		
	Male	Female	Total	Male	Female	Total
Chinese	3722 (73.8%)	515 (39.6%)	4237 (66.8%)	999 (82.6%)	111 (48.9%)	1110 (77.2%)
Non-Chinese	1254 (24.9%)	772 (59.4%)	2026 (31.9%)	211 (17.4%)	116 (51.1%)	327 (22.8%)
Asian	575 (11.4%)	437 (33.6%)	1012 (16.0%)	108 (8.9%)	108 (47.6%)	216 (15.0%)
White	422 (8.4%)	22 (1.7%)	444 (7.0%)	84 (6.9%)	2 (0.9%)	86 (6.0%)
Black	78 (1.5%)	66 (5.1%)	144 (2.3%)	18 (1.5%)	6 (2.6%)	24 (1.7%)
Others	179 (3.5%)	247 (19.0%)	426 (6.7%)	1 (0.1%)	0 (0.0%)	1 (0.1%)
Unknown	67 (1.3%)	12 (0.9%)	79 (1.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Total	5043 (100.0%)	1299 (100.0%)	6342 (100.0%)	1210 (100.0%)	227 (100.0%)	1437 (100.0%)

Box 2.4 Age distribution of reported HIV/AIDS cases


(a) Median age of reported HIV/AIDS cases

Year	HIV			AIDS		
	Median age	Inter quartile range		Median age	Inter quartile range	
		25%	75%		25%	75%
1992	34	28	39.75	39	35.25	44
1993	33	27.5	39	38	30.5	40.5
1994	34	28	40	36	33	40
1995	32	26	40	36	30	44
1996	34	30	41	38	32.25	42.75
1997	35	29	42	37	32	48
1998	34	29	40	39	32	47.5
1999	35	29	43	40	34	51
2000	35	29	43	40	33.5	49.5
2001	34.5	29	42	38	30.75	46.25
2002	36	30	44	41	34	48
2003	36	31	45	39	35	49.25
2004	36	30	44	42	35	51
2005	36	30	44	40	33.75	47.25
2006	34	28	42	38	31	47
2007	34	29	41	41	34	50.5
2008	36	29	45	41	34	54
2009	36	29	44	41	34	51
2010	36	30	44	42	37	53
2011	37	30	47	41	34	48.5
2012	36	29	44	42	35.5	49
2013	36	29	44	43.5	36	49.25
Cumulative (1984 – 2013)	35	29	43	40	33	49


(b) Age & gender of reported HIV cases (Year 2013)




(c) Age & gender of reported AIDS cases (Year 2013)



(d) Age & gender of reported HIV cases (cumulative, 1984 - 2013)



(e) Age & gender of reported AIDS cases (cumulative, 1985 - 2013)



(f) Adults & children with reported HIV/AIDS in 2013

Age	HIV			AIDS		
	Male	Female	Total	Male	Female	Total
Adult	444	114	558	70	13	83
Children (age <=13)	0	1	1	0	1	1
Total	444	115	559	70	14	84

Box 2.5 Exposure category of reported HIV/AIDS case

(a) Distribution of reported HIV cases by exposure category (1994 - 2013)


Exposure Category (%) \ Year	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Cumulative (1984 - 2013)
Heterosexual	73 (70%)	81 (66%)	94 (70%)	117 (65%)	135 (71%)	127 (60%)	115 (63%)	127 (60%)	146 (56%)	117 (51%)	112 (42%)	117 (37%)	130 (35%)	111 (27%)	144 (33%)	116 (29%)	121 (31%)	118 (27%)	134 (26%)	147 (26%)	2523 (40%)
Homosexual	22 (21%)	26 (21%)	20 (15%)	33 (18%)	16 (8%)	34 (16%)	22 (12%)	37 (17%)	48 (18%)	46 (20%)	63 (24%)	87 (28%)	110 (29%)	161 (39%)	139 (32%)	165 (42%)	146 (38%)	178 (41%)	247 (48%)	278 (50%)	2007 (32%)
Bisexual	4 (4%)	4 (3%)	3 (2%)	10 (6%)	6 (3%)	10 (5%)	7 (4%)	7 (3%)	9 (3%)	5 (2%)	6 (2%)	12 (4%)	15 (4%)	19 (5%)	18 (4%)	9 (2%)	23 (6%)	18 (4%)	17 (3%)	22 (4%)	259 (4%)
Injecting drug use	2 (2%)	2 (2%)	1 (1%)	2 (1%)	1 (1%)	6 (3%)	10 (5%)	11 (5%)	10 (4%)	11 (5%)	25 (9%)	31 (10%)	58 (16%)	44 (11%)	42 (10%)	15 (4%)	15 (4%)	14 (3%)	7 (1%)	7 (1%)	323 (5%)
Blood contact	1 (1%)	0 (0%)	0 (0%)	1 (1%)	0 (0%)	2 (1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	4 (1%)	0 (0%)	2 (0%)	3 (1%)	1 (0%)	0 (0%)	2 (0%)	1 (0%)	1 (0%)	83 (1%)
Perinatal	1 (1%)	2 (2%)	1 (1%)	0 (0%)	2 (1%)	4 (2%)	2 (1%)	2 (1%)	1 (0%)	0 (0%)	0 (0%)	2 (1%)	2 (1%)	1 (0%)	0 (0%)	3 (1%)	3 (1%)	0 (0%)	1 (0%)	0 (0%)	27 (0%)
Undetermined	1 (1%)	7 (6%)	15 (11%)	18 (10%)	29 (15%)	30 (14%)	27 (15%)	29 (14%)	46 (18%)	50 (22%)	62 (23%)	60 (19%)	58 (16%)	76 (18%)	89 (20%)	87 (22%)	81 (21%)	108 (25%)	106 (21%)	104 (19%)	1120 (18%)
Total	104 (100%)	122 (100%)	134 (100%)	181 (100%)	189 (100%)	213 (100%)	183 (100%)	213 (100%)	260 (100%)	229 (100%)	268 (100%)	313 (100%)	373 (100%)	414 (100%)	435 (100%)	396 (100%)	389 (100%)	438 (100%)	513 (100%)	559 (100%)	6342 (100%)

(b) Distribution of reported AIDS cases by exposure category (1994 - 2013)


Exposure Category (%) \ Year	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Cumulative (1985 - 2013)
Heterosexual	16 (43%)	31 (69%)	55 (79%)	44 (69%)	50 (79%)	44 (72%)	56 (84%)	49 (82%)	38 (72%)	46 (82%)	35 (71%)	38 (59%)	30 (41%)	40 (51%)	52 (54%)	35 (46%)	36 (46%)	29 (35%)	39 (45%)	31 (37%)	819 (57%)
Homosexual	13 (35%)	9 (20%)	6 (9%)	10 (16%)	6 (10%)	8 (13%)	1 (1%)	5 (8%)	8 (15%)	7 (13%)	8 (16%)	13 (20%)	21 (29%)	20 (25%)	25 (26%)	28 (37%)	27 (34%)	32 (39%)	34 (40%)	36 (43%)	356 (25%)
Bisexual	4 (11%)	3 (7%)	1 (1%)	3 (5%)	1 (2%)	1 (2%)	1 (1%)	2 (3%)	2 (4%)	0 (0%)	0 (0%)	3 (5%)	3 (4%)	1 (1%)	3 (3%)	3 (4%)	5 (6%)	4 (5%)	4 (5%)	5 (6%)	61 (4%)
Injecting drug use	0 (0%)	1 (2%)	1 (1%)	1 (2%)	0 (0%)	1 (2%)	2 (3%)	1 (2%)	1 (2%)	0 (0%)	3 (6%)	1 (2%)	11 (15%)	9 (11%)	9 (9%)	2 (3%)	3 (4%)	5 (6%)	2 (2%)	4 (5%)	59 (4%)
Blood contact	3 (8%)	0 (0%)	2 (3%)	1 (2%)	1 (2%)	2 (3%)	1 (1%)	0 (0%)	0 (0%)	1 (2%)	0 (0%)	1 (2%)	0 (0%)	1 (1%)	2 (2%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	24 (2%)
Perinatal	1 (3%)	1 (2%)	0 (0%)	0 (0%)	1 (2%)	1 (2%)	1 (1%)	1 (2%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (1%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)	8 (1%)
Undetermined	0 (0%)	0 (0%)	5 (7%)	5 (8%)	4 (6%)	4 (7%)	5 (7%)	2 (3%)	4 (8%)	2 (4%)	3 (6%)	8 (13%)	8 (11%)	8 (10%)	5 (5%)	7 (9%)	7 (9%)	12 (15%)	7 (8%)	8 (10%)	110 (8%)
Total	37 (100%)	45 (100%)	70 (100%)	64 (100%)	63 (100%)	61 (100%)	67 (100%)	60 (100%)	53 (100%)	56 (100%)	49 (100%)	64 (100%)	73 (100%)	79 (100%)	96 (100%)	76 (100%)	79 (100%)	82 (100%)	86 (100%)	84 (100%)	1437 (100%)

Box 2.6 Reported HIV/AIDS cases in injecting drug users (1994– 2013)

(a) Reported HIV-infected injecting drug users - by gender




(b) Reported AIDS case in injecting drug users - by gender




Box 2.7 Reported sexually acquired HIV/AIDS cases (1994– 2013)

(a) Yearly reports of sexually acquired HIV cases



(b) Yearly reports of sexually acquired AIDS cases



(c) **Ratio of heterosexual vs. homosexual/bisexual men reported with HIV/AIDS**

Year	HIV	AIDS
1994	2.3 : 1	0.8 : 1
1995	1.9 : 1	2.0 : 1
1996	3.0 : 1	7.1 : 1
1997	2.0 : 1	2.5 : 1
1998	4.2 : 1	5.9 : 1
1999	2.0 : 1	4.2 : 1
2000	2.7 : 1	23.5 : 1
2001	1.9 : 1	5.3 : 1
2002	1.7 : 1	2.7 : 1
2003	1.6 : 1	4.9 : 1
2004	1.1 : 1	3.8 : 1
2005	0.8 : 1	1.8 : 1
2006	0.7 : 1	0.8 : 1
2007	0.4 : 1	1.5 : 1
2008	0.6 : 1	1.4 : 1
2009	0.4 : 1	0.8 : 1
2010	0.4 : 1	0.8 : 1
2011	0.3 : 1	0.4 : 1
2012	0.3 : 1	0.6 : 1
2013	0.2 : 1	0.4 : 1
Cumulative (1984 – 2013)	0.7 : 1	1.5 : 1

Box 2.8 Profile of primary AIDS defining illnesses (ADI) (1994 - 2013)

Year ADI (%)	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	Cumulative (1993 - 2013)
<i>Pneumocystis</i> <i>Pneumonia (PCP)</i>	12 (32%)	17 (38%)	21 (30%)	20 (31%)	26 (41%)	23 (38%)	30 (45%)	26 (43%)	25 (47%)	22 (39%)	22 (45%)	20 (31%)	27 (37%)	28 (35%)	37 (39%)	32 (42%)	36 (46%)	37 (45%)	39 (45%)	37 (44%)	578 (40%)
<i>Mycobacterium</i> <i>Tuberculosis</i>	4 (11%)	8 (18%)	21 (30%)	17 (27%)	18 (29%)	13 (21%)	19 (28%)	17 (28%)	9 (17%)	15 (27%)	13 (27%)	25 (39%)	26 (36%)	32 (41%)	32 (33%)	24 (32%)	20 (25%)	22 (27%)	15 (17%)	17 (20%)	376 (26%)
Other fungal infections	4 (11%)	7 (16%)	6 (9%)	10 (16%)	8 (13%)	5 (8%)	4 (6%)	5 (8%)	8 (15%)	4 (7%)	6 (12%)	5 (8%)	4 (5%)	3 (4%)	3 (3%)	6 (8%)	5 (6%)	8 (10%)	10 (12%)	10 (12%)	132 (9%)
Penicilliosis	6 (16%)	7 (16%)	7 (10%)	5 (8%)	2 (3%)	7 (11%)	5 (7%)	1 (2%)	7 (13%)	5 (9%)	4 (8%)	7 (11%)	11 (15%)	4 (5%)	6 (6%)	1 (1%)	6 (8%)	2 (2%)	6 (7%)	3 (4%)	105 (7%)
Cytomegalovirus diseases	1 (3%)	3 (7%)	4 (6%)	4 (6%)	3 (5%)	2 (3%)	3 (4%)	2 (3%)	0 (0%)	3 (5%)	1 (2%)	2 (3%)	3 (4%)	4 (5%)	6 (6%)	3 (4%)	3 (4%)	5 (6%)	4 (5%)	4 (5%)	66 (5%)
Non-TB mycobacterial infections	0 (0%)	0 (0%)	2 (3%)	1 (2%)	0 (0%)	5 (8%)	1 (1%)	5 (8%)	2 (4%)	1 (2%)	2 (4%)	0 (0%)	1 (1%)	0 (0%)	1 (1%)	2 (3%)	0 (0%)	0 (0%)	2 (2%)	0 (0%)	30 (2%)
Kaposi's sarcoma	4 (11%)	1 (2%)	2 (3%)	3 (5%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (2%)	0 (0%)	1 (2%)	0 (0%)	1 (1%)	4 (4%)	2 (3%)	1 (1%)	2 (2%)	1 (1%)	7 (8%)	37 (3%)
Others	6 (16%)	2 (4%)	7 (10%)	4 (6%)	6 (10%)	6 (10%)	5 (7%)	4 (7%)	2 (4%)	5 (9%)	1 (2%)	4 (6%)	1 (1%)	7 (9%)	7 (7%)	6 (8%)	8 (10%)	6 (7%)	9 (10%)	6 (7%)	113 (8%)
Total	37 (100%)	45 (100%)	70 (100%)	64 (100%)	63 (100%)	61 (100%)	67 (100%)	60 (100%)	53 (100%)	56 (100%)	49 (100%)	64 (100%)	73 (100%)	79 (100%)	96 (100%)	76 (100%)	79 (100%)	82 (100%)	86 (100%)	84 (100%)	1437 (100%)

3. TABULATED RESULTS OF HIV PREVALENCE SURVEYS

System description

- This is a collection of data from HIV prevalence studies and public service records that contribute to the understanding of the HIV situation in selected community groups or settings.

System layout

Target population	Setting	System	Since	Sample size	Data available in 2013
(a) Community with predisposing risk factors					
STI patients	Social Hygiene Clinics	Voluntary testing offered to clients	1985	Around 25000 – 40000 / year	Yes
Drug users (1)	Methadone Clinics	Universal HIV Antibody (Urine samples) Testing Programme	2003	Around 6000 – 9000 / year	Yes
Drug users (2)	Inpatient drug treatment centres/institution	Unlinked anonymous screening (Urine samples)	1998	Around 150 – 700 / year	Yes
Men who have Sex with Men (MSM)	AIDS Concern	Voluntary testing offered to MSM (rapid tests)	2000	Around 200 - 1500 / year	Yes
	HIV Prevalence and Risk behavioural Survey of Men who have sex with men in Hong Kong (PRISM)	Unlinked anonymous screening (urine samples) Voluntary testing (urine samples)	2006 round 2008, 2011 rounds	Around 800 / study	No
Female Sex Worker (FSW)	Community Based Risk Behavioral and Seroprevalence Survey for Female Sex Workers in Hong Kong (CRISP)	Unlinked anonymous screening (urine samples) Voluntary testing (urine samples)	2006 round 2008 round	Around 900/study	No
	HIV and AIDS Response Indicator Survey (HARIS)	Voluntary testing (urine samples)	2013	Around 600/study	Yes
(b) Community without known risk factors					
Blood donors	Hong Kong Red Cross Blood Transfusion Service	A requirement for all potential donors	1985	Around 180000 – 240000 / year	Yes
Antenatal women	All maternal and child health centres and public hospitals	Universal voluntary testing (blood samples)	Sept 2001	Around 40000 - 50000/ year	Yes
(c) Community with undefined risk					
TB patients	TB and Chest Clinics of the Department of Health	Voluntary testing (blood samples)	1993	Around 2000 – 4500 / year	Yes
Prisoners	Penal institutions	Unlinked anonymous screening (blood /urine samples)	1992	Around 1500 – 2500 / year	Yes

Box 3.1 HIV prevalence in blood donors at Hong Kong Red Cross Blood Transfusion Service

(a) HIV detection rate by number of donated blood units (2004 - 2013)

Year	Units of blood donated	No. of units anti-HIV+	Positive detection rate of donated units (%)	95% C.I. for prevalence (%)
2004	198,420	1	0.001	(0.0000 - 0.0028)
2005	197,974	3	0.002	(0.0003 - 0.0044)
2006	196,332	6	0.003	(0.0011 - 0.0067)
2007	205,645	9	0.004	(0.0020 - 0.0083)
2008	212,739	10	0.005	(0.0023 - 0.0086)
2009	214,709	3	0.001	(0.0003 - 0.0041)
2010	224,483	4	0.002	(0.0005 - 0.0046)
2011	234,086	5	0.002	(0.0007 - 0.0050)
2012	241,804	8	0.003	(0.0014 - 0.0065)
2013	244,198	7	0.003	(0.0012 - 0.0059)

(b) HIV prevalence in new and repeat blood donors (2004 - 2013)

Year	New donors			Repeat donors		
	No. of donors	No. of donors anti-HIV+	HIV positivity rate (%) (95% C.I. (%))	No. of donors	No. of donors anti-HIV+	HIV positivity rate (%) (95% C.I. (%))
2004	41,679	0	0 (---)	156,741	1	0.001 (0.0000 – 0.0036)
2005	42,643	1	0.002 (0.0001 – 0.0131)	155,331	2	0.001 (0.0002 – 0.0047)
2006	40,029	2	0.005 (0.0006 – 0.0180)	156,303	4	0.003 (0.0007 – 0.0066)
2007	40,287	3	0.007 (0.0015 – 0.0218)	165,358	6	0.004 (0.0013 – 0.0079)
2008	40,909	5	0.012 (0.0040 – 0.0285)	171,830	5	0.003 (0.0009 – 0.0068)
2009	46,158	1	0.002 (0.0001 – 0.0121)	168,551	2	0.001 (0.0001 – 0.0043)
2010	41,980	2	0.005 (0.0006 – 0.0172)	182,503	2	0.001 (0.0001 – 0.0040)
2011	42,684	2	0.005 (0.0006 – 0.0169)	191,402	3	0.002 (0.0003 – 0.0046)
2012	42,083	3	0.007 (0.0015– 0.0208)	199,721	5	0.003 (0.0008– 0.0058)
2013	40,315	1	0.002 (0.0001– 0.0138)	203,883	6	0.003 (0.0011– 0.0064)

Box 3.2 HIV prevalence in clients attending Social Hygiene Services, from voluntary blood testing (2004 – 2013)

Year	No. of blood samples	No. of samples tested anti-HIV+	Prevalence (%)	95% C.I. for prevalence (%)
2004	43,980	46	0.105	(0.077 - 0.140)
2005	38,978	28	0.072	(0.048 - 0.104)
2006	37,120	47	0.127	(0.093 - 0.168)
2007	33,841	50	0.148	(0.110 - 0.195)
2008	31,040	72	0.232	(0.181 - 0.292)
2009	29,152	50	0.172	(0.127 - 0.226)
2010	26,300	40	0.152	(0.109 - 0.207)
2011	25,599	44	0.172	(0.125 - 0.231)
2012	26,679	55	0.206	(0.155 - 0.268)
2013	26,470	90	0.340	(0.273 - 0.418)

Box 3.3 HIV prevalence in drug users attending methadone clinics

Year	No. of urine samples	No. of samples tested anti-HIV+	Prevalence (%)	95% C.I. for prevalence (%)
2004*	8,812	18	0.204	(0.121 - 0.323)
2005*	8,696	28	0.322	(0.214 - 0.465)
2006*	7,730	28	0.362	(0.241 - 0.524)
2007*	7,314	26	0.355	(0.232 - 0.521)
2008*	7,955	37	0.465	(0.327 - 0.641)
2009*	7,765	38	0.489	(0.346 - 0.672)
2010*	7,445	36	0.484	(0.339 - 0.669)
2011*	6,960	37	0.53	(0.374 - 0.733)
2012*	6,742	42	0.62	(0.449 - 0.842)
Year	Total no. of methadone clinic attendees tested for HIV	Total no. of methadone clinic attendees tested positive for HIV	Prevalence (%)	95% C.I. for prevalence (%)
2013**	6,925	47	0.68	(0.499 - 0.903)

*From the Universal HIV Antibody (Urine) Testing Programme in Methadone clinics

**Overall figures from all methadone clinic attendees

Box 3.4 HIV prevalence in drug users attending inpatient drug treatment centres / institutions, from unlinked anonymous screening (2004 - 2013)

Year	No. of urine samples	No. of samples tested anti-HIV+	Prevalence (%)	95% C.I. for prevalence (%)
2004*	---	---	---	(--- - ---)
2005	630	0	0	(--- - ---)
2006	786	4	0.509	(0.139 - 1.303)
2007	387	0	0	(--- - ---)
2008	369	0	0	(--- - ---)
2009	430	3	0.698	(0.144 - 2.039)
2010	165	0	0	(--- - ---)
2011	396	1	0.253	(0.006 - 1.407)
2012	205	2	0.976	(0.118 - 3.524)
2013	188	0	0	(--- - ---)

* Unlinked anonymous screening was not performed in 2004;

Box 3.5 HIV prevalence in newly admitted prisoners from unlinked anonymous screening (2004 - 2013)

Year	No. of Samples*	No. of samples tested anti-HIV+	Prevalence (%)	95% C.I. for prevalence (%)
2004	1,980	7	0.354	(0.142 - 0.728)
2005	2,007	6	0.299	(0.110 - 0.651)
2006	2,796	13	0.465	(0.248 - 0.795)
2007	2,718	7	0.258	(0.104 - 0.531)
2008	2,231	21	0.941	(0.583 - 1.439)
2009	1,929	15	0.778	(0.435 - 1.283)
2010	1,450	14	0.966	(0.528 - 1.620)
2011	1,445	27	1.869	(1.231 - 2.718)
2012	1,493	11	0.737	(0.368 - 1.318)
2013	1,460	14	0.959	(0.524 - 1.609)

Box 3.6 HIV prevalence in patients attending government TB & Chest Clinics, from voluntary blood testing (2004 - 2013)

Year	No. of blood samples	Coverage*		No. of anti-HIV+	Prevalence (%)	95% C.I. for prevalence (%)
		A	B			
2004	3,202	93.1%	44.4%	10	0.312	(0.150 - 0.574)
2005	4,209	81.2%	68.3%	35	0.832	(0.579 - 1.157)
2006	4,511	91.0%	78.2%	33	0.732	(0.504 - 1.027)
2007	4,075	88.7%	74.6%	41	1.006	(0.722 - 1.365)
2008	4,121	89.9%	73.1%	48	1.165	(0.859 - 1.544)
2009	3,993	89.0%	76.9%	40	1.002	(0.716 - 1.364)
2010	3,833	90.2%	75.3%	28	0.730	(0.485 - 1.056)
2011	3,656	90.6%	76.3%	33	0.903	(0.621 - 1.268)
2012	3,707	91.2%	76.3% [#]	22	0.593	(0.372 - 0.899)
2013	3,536	88.2%	74.1%**	24	0.679	(0.435 - 1.010)

* coverage A is the proportion of patients attended government TB & Chest Clinics who have been tested for HIV in TB Clinic. (For year 2000-2004, it used to be the proportion of patients who started on TB tx at government TB & Chest Clinics who have been tested for HIV in TB Clinic);
 B is the proportion of total TB notifications who have been tested for HIV at government TB & Chest Clinics.

[#] figures revised

** provisional figure

Box 3.7 HIV prevalence among antenatal women from Universal Antenatal HIV Antibody Testing Programme (2004 - 2013)

Year	Number of blood samples	Coverage*	Number of positive tests	Prevalence (%)	95% C.I. for prevalence (%)
2004	41,070	97.9%	6	0.01	(0.0054 - 0.0318)
2005	42,750	98.1%	5	0.01	(0.0038 - 0.0273)
2006	43,297	98.0%	9	0.02	(0.0095 - 0.0395)
2007	47,472	97.4%	11	0.02	(0.0116 - 0.0415)
2008	51,737	98.2%	2	0.004	(0.0005 - 0.0140)
2009	51,227	98.3%	7	0.01	(0.0055 - 0.0282)
2010	55,147	98.6%	10	0.02	(0.0088 - 0.0338)
2011	56,674	98.8%	6	0.01	(0.0039 - 0.0233)
2012	53,857	98.6%	9	0.02	(0.0077 - 0.0322)
2013	49,603	98.5%	7	0.01	(0.0058 - 0.0295)

* coverage is the proportion of women attending public antenatal services who have been tested for HIV

Box 3.8 HIV prevalence among MSM tested by AIDS Concern (2004 - 2013)

Year	Number of test*	Number of positive tests	Prevalence (%)	95% C.I. for prevalence (%)
2004	332	6	1.81	(0.663 - 3.934)
2005	483	12	2.48	(1.284 - 4.340)
2006	610	10	1.64	(0.786 - 3.015)
2007	723	17	2.35	(1.370 - 3.765)
2008	905	15	1.66	(0.928 - 2.734)
2009	909	18	1.98	(1.174 - 3.130)
2010	854	18	2.11	(1.249 - 3.331)
2011	1,026	20	1.95	(1.191 - 3.011)
2012	1,492	30	2.01	(1.357 - 2.871)
2013	1,438	26	1.81	(1.181 - 2.649)

* HIV rapid test

Box 3.9 HIV prevalence among MSM – PRISM* (2006, 2009 and 2011)

Year	Number of urine specimen collected	Number of positive tests	Crude Prevalence (%)	Adjusted Prevalence (%)	95% C.I. for adjusted prevalence (%)
2006	859	37	4.31	4.05	(3.03 - 5.94)
2008	833	37	4.44	4.31	(2.95 - 5.67)
2011	816	30	3.68	4.08	(3.44 - 4.85)

*PRISM: HIV Prevalence and Risk behavioural Survey of Men who have sex with men in Hong Kong, a venue based survey including bars and saunas both in 2006 and 2008 round. Beaches was also added in 2011 round.

Box 3.10 HIV prevalence among Female Sex Workers – CRISP* (2006 and 2009), HARiS ** (2013)

Year	Number of urine specimen collected	Number of positive tests	Adjusted Prevalence (%)
2006	996	5	0.19
2009	986	2	0.05
2013	605	0	0.00%

*CRISP: Community Based Risk Behavioural and Seroprevalence Survey for Female Sex Workers in Hong Kong, a venue based survey including one woman brothels, bars, night clubs, sauna, karaokes etc in 2006 and 2009 round.

**HARiS: HIV and AIDS Response Indicator Survey for Female Sex Workers, a combined venue-based and non-governmental organisations centre-based survey

4. TABULATED RESULTS OF STATISTICS ON SEXUALLY TRANSMITTED INFECTIONS (STI)

System description

- This is a clinic based disease reporting system contributed by Social Hygiene Service, Department of Health. Summary tables are submitted quarterly by Social Hygiene Service. The clinics included in this surveillance system are: Chai Wan, Lek Yuen¹, Wan Chai, Western², Yau Ma Tei, South Kwai Chung³, Yung Fung Shee, Tuen Mun, Fanling ITC⁴, Tai Po, and Shek Wu Hui⁵.

¹Lek Yuen Clinic was closed since April 2005

²Western Social Hygiene Clinic was merged with Wan Chai Social Hygiene Clinic and Sai Ying Pun Dermatology Clinic wef 2.7.2003

³South Kwai Chung Clinic was closed on 27.3.2004

⁴Venereal Diseases Clinics in Fanling ITC was commenced operation in part-time basis on 1.9.2003 by appointment only.


⁵Tai Po and Shek Wu Hui clinics were closed since 2001

Box 4.1 Total number of STI newly reported by individual Social Hygiene Clinic**(a) Year 2013**


	Chai Wan	Wan Chai	Yau Ma Tei	Yung Fung Shee	TuenMun	Fanling ITC [#]	Total
Male	195	1764	3414	1073	610	631	7687
Female	231	986	2453	469	570	516	5225
Total	426	2750	5867	1542	1180	1147	12912

Venereal Diseases Clinics in Fanling ITC commenced operation in part-time basis on 1.9.2003 by appointment only.

(b) 2009 - 2013



Box 4.2 Annual newly reported STIs in Social Hygiene Clinics




^ NSGI / NGU : Non-specific Genital Infection / Non-gonococcal Urethritis

Box 4.3 Syphilis newly reported by Social Hygiene Clinics (2009 - 2013)


Syphilis \ Year	2009	2010	2011	2012	2013
Primary	63	50	52	46	42
Secondary	69	54	51	58	89
Early latent	61	91	64	45	72
Late latent	816	821	805	859	780
Late (cardiovascular / neuro)	12	16	8	3	10
Congenital (early)	0	0	0	0	0
Congenital (late)	3	0	5	2	6
Total	1,024	1,032	985	1,013	999

Box 4.4 Sexually acquired HIV infection in Hong Kong (1994-2013)



* SHS: Social Hygiene Service

Box 4.5 Syndromic presentations of STI from Behavioural Survey of Social Hygiene Service (2009-2013)



5. TABULATED RESULTS ON BEHAVIOURAL MONITORING

System description

- This is a tabulation of HIV risky behavioural data collected from different sources in Hong Kong

System layout

Source	Sexual behaviour	Drug-taking behaviour	Data available in 2013
AIDS Counselling and Testing Service (ACTS), Special Preventive Programme, CHP, DH	<ul style="list-style-type: none"> - Median no. of sexpartners in heterosexual men/MSM - Recent history of commercial sex in heterosexual men - Condom use in heterosexual men/MSM - 		Yes
Social Hygiene Service (SHS)	<ul style="list-style-type: none"> - Recent history of commercial sex / casual sex - Condom use in heterosexual men 		Yes
Methadone clinics (DRS-M)		<ul style="list-style-type: none"> - Proportion of current injectors - Practice of current needle-sharing 	Yes
Shek Kwu Chau (SKC) Treatment and Rehabilitation Centre (DRS-S)		<ul style="list-style-type: none"> - Proportion of current injectors - Practice of current needle-sharing 	Yes
Central Registry of Drug Abuse (CRDA)		<ul style="list-style-type: none"> - Proportion of current injectors in all drug users - Proportion of current injectors in new drug users 	Yes
Street Addict Survey (SAS) (From the Society for the Aid and Rehabilitation of Drug Abusers)		<ul style="list-style-type: none"> - Proportion of current injectors - Practice of current needle-sharing 	Yes
AIDS Concern testing service for MSM (AC)	<ul style="list-style-type: none"> - Condom use in MSM 		Yes
HIV Prevalence and Risk behavioural Survey of Men who have sex with men in Hong Kong (PRISM)	<ul style="list-style-type: none"> - Condom use in MSM 		No
HIV and AIDS Response Indicator Survey (HARiS)	<ul style="list-style-type: none"> - Condom use in MSM 		Yes

Box 5.1 Median number of sex partners in the previous year among adult[^] heterosexual men / MSM attending AIDS Counselling and Testing Service (ACTS) (2004-2013)

	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Heterosexual men - Regular sex partners [*]	1	1	1	1	1	1	1	1	1	1
Heterosexual men - Commercial sex partners ^{**}	2	2	2	2	2	3	3	2	3	2
Heterosexual men - Casual sex partners ^{***}	1	1	1	1	1	1	1	1	1	1
MSM - Regular sex partners [*]	1	1	1	1	1	1	1	1	1	1
MSM - Commercial sex partners ^{**}	2	1	1.5	1	2	3	1.5	1	2	4.5
MSM - Casual sex partners ^{***}	4	3	3	3	4	4	3.5	3	3	3


[^] Adult: aged 18 or above

^{*} Regular sex partners used to refer to long-term sex partners including spouse, mistress, and steady boyfriends/girlfriends for at least one year, or if less than one year, one with whom is expected to continue sexual relationship. This definition of regular sex partners in 2008 has been further refined to include (other than the long-term sex partners) sex buddy that refers to regular sex only partner for at least 6 months, or if less than 6 months, one with whom is expected to continue sexual relationship

^{**} Commercial sex partners are defined as those who have sexual intercourse in exchange for money, goods or services. Examples are prostitutes and customers of prostitutes.

^{***} Casual sex partners, the two do not have steady relationship.

Box 5.2 Recent history* of commercial / casual sex among adult^ heterosexual men (2004-2013)



* Commercial sex partners are defined as those who have sexual intercourse in exchange for money, goods or services. Examples are female sex workers and their clients. Casual sex partners are defined as those who are non-regular and non-commercial. Examples are those on one-night stand. SHS & ACTS refers to such history in past one year;


^ Adult: aged 18 or above

Remarks : SHS – Social Hygiene Services

ACTS - AIDS Counselling and Testing Service

Box 5.3 Condom use with regular partners among adult heterosexual men

(a) Consistent condom use* with regular partners** among adult^ heterosexual men (2004-2013)



* Consistent condom use is defined as always or 100% of the time using a condom


ACTS captures such condom usage in past one year while SHS captures such usage in past 3 months

** Regular sex partners used to refer to long-term sex partners including spouse, mistress, and steady girl friends for at least one year, or if less than one year, one with whom is expected to continue sexual relationship. This definition of regular sex partners in 2008 has been further refined to include (other than the long-term sex partners) sex buddy that refers to regular sex only partner for at least 6 months, or if less than 6 months, one with whom is expected to continue sexual relationship

^ Adult: aged 18 or above

Remarks : SHS – Social Hygiene Services, ACTS - AIDS Counselling and Testing Service

(b) Condom use for last sex with regular partners* among adult^ heterosexual men (2004-2013)



* Regular sex partners used to refer to long-term sex partners including spouse, mistress, and steady girl friends for at least one year, or if less than one year, one with whom is expected to continue sexual relationship. This definition of regular sex partners in 2008 has been further refined to include (other than the long-term sex partners) sex buddy that refers to regular sex only partner for at least 6 months, or if less than 6 months, one with whom is expected to continue sexual relationship


^ Adult: aged 18 or above

Remarks : SHS – Social Hygiene Services

ACTS - AIDS Counselling and Testing Service

Box 5.4 Condom use with commercial / casual partners among adult heterosexual men

(a) Consistent condom use* with commercial / casual partners** among adult^ heterosexual men (2004-2013)



* Consistent condom use is defined as always or 100% of the time using a condom


ACTS captures such condom usage in past one year while SHS captures such usage in past 3 months

** Commercial sex partners are defined as those who have sexual intercourse in exchange for money, goods or services. Examples are female sex workers and their clients. Casual sex partners are defined as those who are non-regular and non-commercial. Examples are those on one-night stand.

^ Adult: aged 18 or above

Remarks : SHS – Social Hygiene Services
ACTS - AIDS Counselling and Testing Service

(b) Condom use for last sex with commercial / casual partners* among adult^ heterosexual men (2004-2013)




* Commercial sex partners are defined as those who have sexual intercourse in exchange for money, goods or services. Examples are female sex workers and their clients. Casual sex partners are defined as those who are non-regular and non-commercial. Examples are those on one-night stand.

^ Adult: aged 18 or above

Remarks : SHS – Social Hygiene Services, ACTS - AIDS Counselling and Testing Service

Box 5.5 Condom use among Men have Sex with Men (MSM)

(a) Consistent condom use* among MSM (2004-2013)



* Consistent condom use is defined as always or 100% of the time using a condom. ACTS captures such condom usage in past one year while AC captures such usage in past 3 months

** Regular sex partners used to refer to long-term sex partners including spouse, mistress, and steady boy/girl friends for at least one year, or if less than one year, one with whom is expected to continue sexual relationship. This definition of regular sex partners in 2008 has been further refined to include (other than the long-term sex partners) sex buddy that refers to regular sex only partner for at least 6 months, or if less than 6 months, one with whom is expected to continue sexual relationship

*** Casual sex partners, the two do not have steady relationship.


**** The data in 2012 only from January to March

Since April 2012, the sex partner types from AC survey further breakdown into regular sex partner, casual sex partner and boyfriend

^ Adult: aged 18 or above


Remarks: ACTS - AIDS Counselling and Testing Service, AC - AIDS Concern, PRISM- HIV Prevalence and Risk Behavioural Survey of MSM in Hong Kong

(b) Condom use for last anal sex among MSM (2004-2013)



- * Regular sex partners used to refer to long-term sex partners including spouse, and steady boy friends for at least one year, or if less than one year, one with whom is expected to continue sexual relationship. This definition of regular sex partners in 2008 has been further refined to include (other than the long-term sex partners) sex buddy that refers to regular sex only partner for at least 6 months, or if less than 6 months, one with whom is expected to continue sexual relationship
- ** Casual sex partners, the two do not have steady relationship.
- *** The data in 2012 only from January to March
- ^ Adult: aged 18 or above
- # Since April 2012, the sex partner types from AC survey further breakdown into regular sex partner, casual sex partner and boyfriend
- Remarks : ACTS - AIDS Counselling and Testing Service
AC - AIDS Concern, HARiS - HIV and AIDS Response Indicator Survey

Box 5.6 Proportion of current injectors* (2004-2013)




* Definitions differ for different data sources. DRS-S refers to drug injecting behaviour in past 6 months (before 2006, it referred to drug injecting at the time of programme admission); DRS-M refers to drug injecting at the time of programme admission; SAS refers to drug injecting behaviour in past 1 month (before 2007, it referred to drug injecting in past 3 months); CRDA refers to drug injecting behaviour in past 4 weeks;

** New clients refer to people who are known to the CRDA for the first time in a period. For a particular period, a person will be regarded as a newly reported person if and only if the person does not have any report before the specified period.

Remarks: DRS-S - Shek Kwu Chau Treatment and Rehabilitation Centre (Newly / Re-admitted case)
 DRS-M - Methadone clinics (Newly admitted case only)
 SAS - Street Addict Survey (From the Society for the Aid and Rehabilitation of Drug Abusers (SARDA))
 CRDA - Central Registry of Drug Abuse

Box 5.7 Proportion of current needle-sharers* (2004-2013)



* This figure referred to the proportion of current syringe sharing behaviour among current injectors. Definitions differ for different data sources. DRS-S refers to such sharing behaviour among those who injected drug in past 6 months (before 2006, it referred to such sharing behaviour in past 6 months among those who injected drug at the time of programme admission); SAS refers to such sharing behaviour among those who injected drug in past 1 month (before 2007, it referred to such sharing behaviour in past 3 months); DRS-M refers to such sharing behaviour in past 4 weeks among those who injected drug at the time of programme admission;


Remarks: DRS-S - Shek Kwu Chau Treatment and Rehabilitation Centre (Newly / Re-admitted cases)
 DRS-M - Methadone clinics (Newly admitted case only)
 SAS - Street Addict Survey (From the Society for the Aid and Rehabilitation of Drug Abusers (SARDA))

6. TABULATED RESULTS OF HIV-1 GENOTYPING STUDIES

System description

- This is a laboratory based reporting system contributed by Virology Division of Public Health Laboratory Services Branch, Centre for Health Protection, Department of Health. HIV viral isolates are collected from the confirmatory laboratories for subtype analysis which are collated with epidemiological information when available. Subtype results are submitted monthly by Virology Division. The confirmatory laboratories included in this surveillance system are: DH Public Health Laboratory Service Branch, Microbiology laboratories of Queen Elizabeth Hospital, Prince of Wales Hospital, Hong Kong Red Cross Blood Transfusion Service. Subtype analysis was commenced since 2001


Box 6.1 Proportion of reports* with subtypes by year in Hong Kong, 2001 - 2013




*: including cases with HIV type 1 or PCR positive result.

Box 6.2 Distribution of HIV-1* subtypes

(i) Cumulative (2001-2013)




(ii) Year 2013



*: including cases with HIV type 1 or PCR positive result.

**: including subtype A, A1, A2, B', D, F, F1, G, CRF03_AB, CRF05_DF, CRF06_CPX, CRF10_CD, CRF11_CPX, CRF12_BF, CRF13_cpx, CRF14_BG, CRF15_01B, CRF55_01B.


Box 6.3 Trend in the common HIV-1* subtypes in Hong Kong, 2001 – 2013



*: including cases with HIV type 1 or PCR positive result.


Box 6.4 Trend in HIV-1* subtype CRF01_AE in Hong Kong, 2001 – 2013

(a) By gender (proportion of cases with subtype CRF01_AE)




*: including cases with HIV type 1 or PCR positive result.

(b) By ethnicity (proportion of cases with subtype CRF01_AE)




(c) By route of transmission (proportion of cases with subtype CRF01_AE)




Box 6.5 Trend in HIV-1* subtype B in Hong Kong, 2001 – 2013

(a) By gender (proportion of cases with subtype B)




*: including cases with HIV type 1 or PCR positive result.

(b) By ethnicity (proportion of cases with subtype B)




(c) By route of transmission (proportion of cases with subtype B)




Box 6.6 Trend in HIV-1* subtype C in Hong Kong, 2001 – 2013

(a) By gender (proportion of cases with subtype C)




*: including cases with HIV type 1 or PCR positive result.


(b) By ethnicity (proportion of cases with subtype C)



(c) By route of transmission (proportion of cases with subtype C)



Box 6.7 Prevalence of intermediate or high level drug resistance related mutation among newly diagnosed HIV patients, 2003-2012



Appendix I: HIV/AIDS report form (DH2293)

DEPARTMENT OF HEALTH
HIV/AIDS Report Form

The HIV/AIDS voluntary reporting system has been in place since 1984. All doctors are encouraged to report patients with HIV/AIDS and to update status of the previously reported cases where appropriate. This is an anonymous and confidential system. Data collected is crucial for understanding the HIV epidemiology in Hong Kong and is used in global analysis only. Aggregate statistics are released quarterly and can be obtained at <http://www.aids.gov.hk>. For any query, please call 3143 7225 or email us at aids@dh.gov.hk

Please complete ALL sections and '✓' in the appropriate box.

Section (A) – Report of HIV

[1] THIS is a ☐ NEW report or ☐ UPDATE of previous reported case

[2] Your reference code numberⁱ: [3] Does the patient have a HK identity card? ☐ Yes ☐ No

[4] Sex : ☐ M ☐ F For female, is she pregnant? ☐ No ☐ Yes If yes, go to **Box I**

[5] Date of birth:/...../..... (ddmmmyyy) OR Age at last birthday:

[6] Ethnicity: ☐ Chinese ☐ Asian ☐ Caucasian ☐ Black ☐ Others: ☐ Unknown

[7] Suspected risk(s) for HIV infectionⁱⁱ

☐ Heterosexual ☐ Homosexual ☐ Bisexual

☐ Injecting drug use

☐ Transfusion of blood/blood products (Haemophilia: ☐ Yes ☐ No)

☐ Perinatal

☐ Others, please specify:

☐ Asked, but risk undetermined

☐ Not asked

Box I

Gravida..... Para..... LMP:/...../..... (ddmmmyyy)

Obstetric follow up clinic/ hospital :

Plan: ☐ TOP ☐ Continue pregnancy

Expected hospital/place of delivery:

[8] Suspected place of infection: ☐ Hong Kong ☐ Mainland China, specify: ☐ Others, specify:
☐ Asked, but undetermined ☐ Not asked

[9] Date of laboratory diagnosis in HK:/...../..... (ddmmmyyy) [10] Western blot confirmation: ☐ Yes ☐ No

[11] Name of Laboratory: [12] Laboratory Number, if a/v:

[13] Previous HIV diagnosis outside HK: ☐ No ☐ Yes If yes, date:/...../..... (ddmmmyyy) place:

[14] Date of last negative HIV test:/...../..... (ddmmmyyy)

[15] CD4 (cells/μl): Date:/...../..... (ddmmmyyy)

[16] HIV status of spouse/regular partner: ☐ HIV positive ☐ HIV negative ☐ Unknown

Section (B) – Report of AIDS

[17] Has the patient developed AIDSⁱⁱⁱ: ☐ Yes ☐ No (Go to Section C)

[18] If yes, the AIDS defining illness(es) is (are):

(i) Date of diagnosis:/...../..... (ddmmmyyy)

(ii) Date of diagnosis:/...../..... (ddmmmyyy)

(iii) Date of diagnosis:/...../..... (ddmmmyyy)

[19] CD4 (cells/μl) at AIDS: Date:/...../..... (ddmmmyyy)

Section (C) – Report of deaths and defaults

[20] Has the patient died? ☐ Yes ☐ No If yes, date of death:/...../..... (ddmmmyyy) Cause:

[21] Has the patient left HK/defaulted follow up? ☐ Yes ☐ No If yes, last seen on:/...../..... (ddmmmyyy)

Section (D) – Correspondence

Name of medical practitioner: ☐ in private practice ☐ in public service

Correspondence Address:

Tel: Fax:

Email: Date:/...../..... (ddmmmyyy)

ⁱ Please put down any code of your choice (e.g., case number) for matching purpose only.


ⁱⁱ Please tick the most likely risk for contracting HIV infection. If there is more than 1 suspected risks, please put down 1 & 2 in descending order of the two most likely risks

ⁱⁱⁱ Surveillance definition of AIDS: a definitive laboratory diagnosis of HIV infection AND one or more of the AIDS indicator conditions (July 1995, Scientific Committee on AIDS. Available at: <http://www.aids.gov.hk/report.htm>).

Appendix II: Classification system for HIV infection and surveillance case definition for AIDS in adolescents and adults in Hong Kong.

A definitive laboratory diagnosis of HIV infection normally by a positive screening test for HIV antibody (e.g. ELISA) supplemented by a confirmatory test (e.g. western blot) + one or more of the AIDS indicator conditions	
AIDS indicator conditions	Candidiasis of bronchi, trachea, or lungs Candidiasis, oesophageal Cervical cancer, invasive Coccidioidomycosis, disseminated or extrapulmonary Cryptococcosis, extrapulmonary Cryptosporidiosis, chronic intestinal (>1 month's duration) Cytomegalovirus disease (other than liver, spleen or nodes) Cytomegalovirus retinitis (with loss of vision) Encephalopathy, HIV-related <i>Herpes simplex</i> : chronic ulcer(s) (>1 month's duration); or bronchitis, pneumonitis, or oesophagitis Histoplasmosis, disseminated or extrapulmonary Isosporiasis, chronic intestinal (>1 month's duration) Kaposi's sarcoma Lymphoma, Burkitt's (or equivalent term) Lymphoma, primary, of brain <i>Mycobacterium tuberculosis</i> ; extrapulmonary or pulmonary/cervical lymph node (only if CD4 < 200/ul) Pneumonia, recurrent Penicilliosis, disseminated Mycobacterium, other species or unidentified species, disseminated or extrapulmonary <i>Pneumocystis carinii</i> pneumonia Progressive multifocal leukoencephalopathy Salmonella septicaemia, recurrent Toxoplasmosis of brain Wasting syndrome due to HIV
Hong Kong has adopted the 1993 Centers for Disease Control and Prevention (CDC) AIDS classification with 3 modifications: (1) disseminated penicilliosis is added as one AIDS-defining condition, (2) pulmonary or cervical lymph node tuberculosis included only if CD4 < 200 µl, (3) a CD4 < 200 µl without any AIDS-defining condition is not counted as AIDS.	

Appendix III: Condom distribution for the prevention of HIV and STI by Department of Health



Note:

1. Community network includes collaborative projects with Action for REACH OUT, AIDS Concern, CHOICE and Phoenix Project of SARDA
2. AIDS Unit and others condom distribution points, such as Travel Health Centres, Correctional Service Department and Tuberculosis and Chest Clinic (started in October 2012).